arXiv:1702.01150v1 [math.GT] 3 Feb 2017

GENERATING SETS OF REIDEMEISTER MOVES OF ORIENTED SINGULAR LINKS AND QUANDLES

KHALED BATAINEH, MOHAMED ELHAMDADI, MUSTAFA HAJIJ, AND WILLIAM YOUMANS

Abstract. We give a generating set of the generalized Reidemeister moves for oriented singular links. We use it to introduce an algebraic structure arising from the study of oriented singular knots. We give some examples, including some non-isomorphic families of such structures over non-abelian groups. We show that the set of colorings of a singular knot by this new structure is an invariant of oriented singular knots and use it to distinguish some singular links.

Contents

1. Introduction 1
2. Basics of quandles 2
3. Oriented singular knots and quandles 2
4. Oriented singquandles over groups 6
5. A generating set of oriented singular Reidemeister moves 8
6. Open questions 13
References 13

1. Introduction

The discovery of the Jones polynomial of links [10] generated a search which uncovered vast families of invariants of knots and links, among them the Vassiliev knot invariants [16]. The Jones polynomial and its relatives can be computed combinatorially using knot diagrams or their braid representations. A fundamental relationship between the Jones polynomial and Vassiliev invariants was established in the work of Birman and Lin [4] where they showed that Vassiliev invariants can be characterized by three axioms. Instead of focusing on a given knot, Vassiliev changed the classical approach by deciding to study the space of all knots instead. This allowed one to study the space of all knots instead of focusing on a single knot. As a result of this work, Vassiliev generated the theory of singular knots and their invariants and has gained considerable attention since then.

Singular knots and their invariants have proven to be important subjects of study on their own, and many classical knot invariants have been successfully extended to singular knots. For example, the work of Fiedler [9] where the author extended the Jones and Alexander polynomials to singular knots. The colored Jones polynomial was generalized to singular knots in [2]. Jones-type invariants for singular links were constructed using a Markov trace on a version of Hecke algebras in [12].

The main purpose of this paper is to relate the theory of quandles to theory of singular knots. In [6], the authors developed a type of involutory quandle structure called *singquandles* to study non-oriented singular knots. This article solves the last open question given in [6] by introducing certain algebraic structures with the intent of applying them to the case of oriented singular knots.

2000 *Mathematics Subject Classification.* Primary 57M25.
Key words and phrases. Generating sets of Reidemeister moves; Quandles, Singular Links.

We call these structures *oriented singquandles*. We give multiple examples of such structures and use them to distinguish between various singular knots. Finally, for the purpose of constructing the axioms of singquandles, it was necessary to construct a generating set of Reidemeister moves acting on oriented singular links which can be found in section 5.

Organization. This article is organized as follows. In section 2 we review the basics of quandles. In section 3 we introduce the notion of oriented singquandles. In section 4 we focus our work on oriented singquandles whose underlying structures rely on group actions and provide some applications of singquandles to singular knot theory. Finally, in section 5 we detail a generating set of Reidemeister moves for oriented singular knots and links.

2. Basics of quandles

Before we introduce any algebraic structures related to singular oriented links, we will need to recall the definition of a quandle and give a few examples. For a more detailed exposition on quandles, see [7, 11, 13].

Definition 2.1. A *quandle* is a set X with a binary operation $(a, b) \mapsto a * b$ such that the following axioms hold:

(1) For any $a \in X$, $a * a = a$.
(2) For any $a, b \in X$, there is a unique $x \in X$ such that $a = x * b$.
(3) For any $a, b, c \in X$, we have $(a * b) * c = (a * c) * (b * c)$.

Axiom (2) of Definition 2.1 states that for each $y \in X$, the map $*_y : X \to X$ with $*_y(x) := x * y$ is a bijection. Its inverse will be denoted by the mapping $\overline{*}_y : X \to X$ with $\overline{*}_y(x) = x \ \overline{*} \ y$, so that $(x * y) \ \overline{*} \ y = x = (x \ \overline{*} \ y) * y$. Below we provide some typical examples of quandles.

- Any set X with the operation $x * y = x$ for any $x, y \in X$ is a quandle called the *trivial* quandle.
- A group $X = G$ with n-fold conjugation as the quandle operation: $x * y = y^{-n}xy^n$.
- Let n be a positive integer. For elements $x, y \in \mathbb{Z}_n$ (integers modulo n), define $x * y \equiv 2y - x \pmod{n}$. Then $*$ defines a quandle structure called the *dihedral quandle*, R_n.
- For any $\mathbb{Z}[T, T^{-1}]$-module M, $x * y = Tx + (1 - T)x$ where $x, y \in M$ defines an *Alexander quandle*.
- Let $< , >: \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ be a symmetric bilinear form on $\mathbb{R}^n$. Let X be the subset of $\mathbb{R}^n$ consisting of vectors x such that $< x, x > \neq 0$. Then the operation
$$x * y = \frac{2 < x, y >}{< x, x >} y - x$$
defines a quandle structure on X. Note that, $x * y$ is the image of x under the reflection in y. This quandle is called a *Coxeter* quandle.

A function $\phi : (X, *) \to (Y, \diamond)$ is a quandle *homomorphism* if $\phi(a * b) = \phi(a) \diamond \phi(b)$ for any $a, b \in X$. A bijective quandle endomorphism of $(X, *)$ is called a quandle isomorphism. For example any map $f_{a,b} : (\mathbb{Z}_n, *) \to (\mathbb{Z}_n, *)$ defined by $f_{a,b}(x) = ax + b$ with $a, b \in \mathbb{Z}_n$ and a invertible in $\mathbb{Z}_n$ is a quandle isomorphism, where $x * y = 2y - x$ (see [8] for more details).

3. Oriented singular knots and quandles

Recall that a singular link in S^3 is the image of a smooth immersion of n circles S^3 that has finitely many double points, called singular points. An orientation of each circle induces orientation on each component of the link. This given an oriented singular link. In this paper we will assume that any singular link is oriented unless specified otherwise. Furthermore, we will work with singular link projections, or diagrams, which are projections of the singular link to the plane such that the information at each crossing is preserved by leaving a little break in the lower strand. Two oriented

singular link diagrams are considered equivalent if and only if one can obtain one from the other by a finite sequence of singular Reidemeister moves (see Figure 2).

In the case of classical knot theory, the axiomatization of the Reidemeister moves gives rise to the definition of a quandle. One of the goals of this paper is to generalize the structure of quandles by considering singular oriented knots and links. We will call this structure an *oriented singquandle* (see the definition below). The axioms of oriented singquandles will be constructed using a generating set of Reidemeister moves on oriented singular links (see Figure 2).

A semiarc in a singular link diagram L is an edge in the link L considered as a 4-valent graph. The oriented singquandle axioms are obtained by associating elements of the oriented singquandle to semiarcs in an oriented singular link diagram and letting these elements act on each other at crossings as shown in the following figure:



FIGURE 1. Regular and singular crossings

Now the goal is to derive the axioms that the binary operators R_1 and R_2 in Figure 1 should satisfy. For this purpose, we begin with the generating set of Reidemeister moves given in Figure 2. The proof that this is a generating set will be postponed to section 5. Using this set of Reidemeister moves, the singquandle axioms can be derived easily. This can be seen in Figures 3, 4 and 5.



FIGURE 3. The Reidemeister move $\Omega 4a$ and colorings



FIGURE 2. A generating set of singular Reidemeister moves



FIGURE 4. The Reidemeister move Ω4e and colorings



FIGURE 5. The Reidemeister move $\Omega 5a$ and colorings

The previous three figures immediately gives us the following definition.

Definition 3.1. Let $(X, *)$ be a quandle. Let R_1 and R_2 be two maps from $X \times X$ to X. The triple $(X, *, R_1, R_2)$ is called an *oriented singquandle* if the following axioms are satisfied:

$$R_1(x \bar{*} y, z) * y = R_1(x, z * y) \quad \text{coming from } \Omega 4\text{a} \tag{3.1}$$

$$R_2(x \bar{*} y, z) = R_2(x, z * y) \bar{*} y \quad \text{coming from } \Omega 4\text{a} \tag{3.2}$$

$$(y \bar{*} R_1(x, z)) * x = (y * R_2(x, z)) \bar{*} z \quad \text{coming from } \Omega 4\text{e} \tag{3.3}$$

$$R_2(x, y) = R_1(y, x * y) \quad \text{coming from } \Omega 5\text{a} \tag{3.4}$$

$$R_1(x, y) * R_2(x, y) = R_2(y, x * y) \quad \text{coming from } \Omega 5\text{a} \tag{3.5}$$

We give the following examples.

Example 3.2. Let $x * y = ax + (1 - a)y$, where a is invertible so that $x \bar{*} y = a^{-1}x + (1 - a^{-1})y$. Now let $R_1(x, y) = bx + cy$, then by axiom 2.4 we have $R_2(x, y) = acx + (c(1-a)+b)y$. By plugging these expressions into the above axioms we can find the relation $c = 1 - b$. Substituting, we find that the following is an oriented singquandle for any invertible a and any b in $\mathbb{Z}_n$:

$$x * y = ax + (1 - a)y \tag{3.6}$$

$$R_1(x, y) = bx + (1 - b)y \tag{3.7}$$

$$R_2(x, y) = a(1 - b)x + (1 - a(1 - b))y \tag{3.8}$$

It is worth noting that all of the above relations between constants can be derived from axiom 2.3, and the other axioms provide only trivial identities or the same relations. In this way, we can extend this generalized affine singquandle to the nonoriented case as well by allowing $x \bar{*} y = x * y$, since axiom 2.3 will reduce to it's counterpart in the axioms of nonoriented singquandles. (See axiom 4.1 given in [6]).

With this observation we can generalize the class of involutive Alexander quandles into a class of singquandles, which is given in the following example.

Example 3.3. Let $\Lambda = \mathbb{Z}[t^{\pm 1}, v]$ and let X be a Λ-module. Then the operations

$$x * y = tx + (1 - t)y, \quad R_1(x, y) = \alpha(a, b, c)x + (1 - \alpha(a, b, c))y \quad \text{and}$$

$$R_2(x, y) = t(1 - \alpha(a, b, c))x + (1 - t)(1 - \alpha(a, b, c))y$$

where $\alpha(a, b, c) = at + bv + ctv$, make X an oriented singuandle we call an *Alexander oriented singquandle*. That X is an oriented singquandle follows from proposition 3.2 by straightforward substitution.

A *coloring* of an oriented singular link L is a function $C : R \longrightarrow L$, where X is a fixed oriented singquandle and R is the set of semiarcs in a fixed diagram of L, satisfying the conditions given in Figure 1. Now the following lemma is immediate from Definition 3.1.

Lemma 3.4. The set of colorings of a singular knot by an oriented singquandle is an invariant of oriented singular knots.

The set of colorings of a singular link L by an oriented singquandle X will be denoted by $Col_X(L)$. As in the usual context of quandles, the notions of oriented singquandle homomorphisms and isomorphisms are immediate.

Definition 3.5. Let $(X, *, R_1, R_2)$ and $(Y, \triangleright, S_1, S_2)$ be two oriented singquandles. A map $f : X \longrightarrow Y$ is homomorphism if the following axioms are satisfied :

(1) $f(x * y) = f(x) \triangleright f(y)$,
(2) $f(R_1(x, y)) = S_1(f(x), f(y))$,
(3) $f(R_2(x, y)) = S_2(f(x), f(y))$.

If in addition f is a bijection then we say that $(X, *, R_1, R_2)$ and $(Y, \triangleright, S_1, S_2)$ are isomorphic. Note that isomorphic oriented singquandles will induce the same set of colorings.

4. Oriented singquandles over groups

When the underlying set of an oriented singquandle X is a group one obtains a rich family of structures. In this section we give a variety of examples including a generalization of affine oriented singquandles as well as an infinite family of non-isomophic singquandles over groups.

Example 4.1. Let $X = G$ be an abelian group, with f being a group automorphism and g a group endomorphism. Consider the operations $x * y = f(x) + y - f(y)$ and $R_1(x, y) = g(y) + x - g(x)$. We can deduce from axiom 2.4 that $R_2(x, y) = g(f(x)) + y - g(f(y))$. Plugging into the axioms, we find that the axioms are satisfied when $(f \circ g)(x) = (g \circ f)(x)$. This structure will reappear in example 3.2, which follows as a special case where $f(x) = ax$ and $g(x) = (1 - b)x$.

Example 4.2. Let $X = G$ be a group, and take the n-fold conjugation quandle from Definition 2.1 with $n = 1$ such that $x * y = y^{-1}xy$. Then a direct computation gives the fact that $(X, *, R_1, R_2)$ is a singquandle if and only if R_1 and R_2 satisfy the following equations:

$$y^{-1}R_1(yxy^{-1}, z)y = R_1(x, y^{-1}zy) \tag{4.1}$$
$$R_2(yxy^{-1}, z) = yR_2(x, y^{-1}zy)y^{-1} \tag{4.2}$$
$$x^{-1}R_1(x, z)y[R_1(x, z)]^{-1}x = z[R_2(x, z)]^{-1}y[R_2(x, z)]z^{-1} \tag{4.3}$$
$$R_2(x, y) = R_1(y, y^{-1}xy) \tag{4.4}$$
$$[R_2(x, y)]^{-1} R_1(x, y)R_2(x, y) = R_2(y, y^{-1}xy) \tag{4.5}$$

A straightforward computation gives the following solutions, for all $x, y \in G$.

(1) $R_1(x, y) = x$ and $R_2(x, y) = y$.
(2) $R_1(x, y) = xyxy^{-1}x^{-1}$ and $R_2(x, y) = xyx^{-1}$.
(3) $R_1(x, y) = y^{-1}xy$ and $R_2(x, y) = y^{-1}x^{-1}yxy$.
(4) $R_1(x, y) = xy^{-1}x^{-1}yx$, and $R_2(x, y) = x^{-1}y^{-1}xy^2$.
(5) $R_1(x, y) = y(x^{-1}y)^n$ and $R_2(x, y) = (y^{-1}x)^{n+1}y$, where $n \geq 1$.

Next we focus our attention on a subset of infinite families of oriented singquandle structures in order to show that some in fact are not isomorphic.

Proposition 4.3. Let $X = G$ be a non-abelian group with the binary operation $x * y = y^{-1}xy$. Then, for $n \geq 1$, the following families of maps R_1 and R_2 give pairwise non-isomorphic oriented singquandles structures $(X, *, R_1, R_2)$ is an oriented singquandle on G:

(1) $R_1(x, y) = x(xy^{-1})^n$ and $R_2(x, y) = y(x^{-1}y)^n$,
(2) $R_1(x, y) = (xy^{-1})^n x$ and $R_2(x, y) = (x^{-1}y)^n y$,
(3) $R_1(x, y) = x(yx^{-1})^{n+1}$ and $R_2(x, y) = x(y^{-1}x)^n$.

Furthermore, in each of the cases (1), (2) and (3), different values of n give also non-isomorphic singular quandles.

Proof. A direct computations show that R_1 and R_2 satisfy the five axioms of Definition 3.1.



FIGURE 6. An oriented Hopf link

To see that the three solutions are pairwise non-isomorphic singquandles, we compute the set of colorings of a singular knot by each of the three solutions. Consider the singular link given in figure 6 and color the top arcs by elements x and y of G. Then it is easy to see that the set of colorings is given by

$$Col_X(L) = \{(x, y) \in G \times G \mid x = R_1(y, y^{-1}xy)\ y = R_2(y, y^{-1}xy)\}. \tag{4.6}$$

Using $R_1(x, y) = x(xy^{-1})^n$ and $R_2(x, y) = y(x^{-1}y)^n$ from solution (1), the set of colorings becomes:

$$Col_X(L) = \{(x, y) \in G \times G \mid (x^{-1}y)^{n+1} = 1\}, \tag{4.7}$$

while the set of colorings of the link L with $R_1(x, y) = (xy^{-1})^n x$ and $R_2(x, y) = (x^{-1}y)^n y$ from solution (2) is:

$$Col_X(L) = \{(x, y) \in G \times G \mid x^{-1}(x^{-1}y)^n y = 1\}. \tag{4.8}$$

Finally, the set of colorings of the same link with $R_1(x, y) = x(yx^{-1})^{n+1}$ and $R_2(x, y) = x(y^{-1}x)^n$ from solution (3) is:

$$Col_X(L) = \{(x, y) \in G \times G \mid (y^{-1}x)^n = 1\}. \tag{4.9}$$

This allows us to conclude that solutions (1), (2), and (3) are pairwise non-isomorphic oriented singquandles. In fact these computations also give that different values of n in any of the three solutions (1), (2) and (3) give non-isomorphic oriented singquandles. We exclude the case of $n = 0$ as solutions (1) and (2) become equivalent.

□

We illustrate here how these new structures can be used to distinguish between oriented singular knots and links.

Example 4.4. We choose X to be a non-abelian group and consider the 1-fold conjugation quandle on X along with the binary operation R_1 and R_2 given by $R_1(x, y) = x^2y^{-1}$ and $R_2(x, y) = yx^{-1}y$. By considering solution (5) in Example 4.2 when $n = 1$ we know that X is an oriented singquandle. In this example we show how the this oriented singquandle can be used to distinguish between two singular knots that differ only in orientation.

Color the two arcs on the top of the knot on the left of Figure 7 by elements x and y. This implies that the coloring space is the set in the diagonal in $G \times G$. On the other hand, the knot on the righthand side of Figure 7 has the coloring space $\{(x, y) \in G \times G,\ \ xyx^{-1} = yxy^{-1}\}$. Since this

set is not the same as the diagonal of $G \times G$, the coloring invariant distinguishes these two oriented singular knots.

7



FIGURE 7. Two oriented singular Hopf links with one singular point

5. A generating set of oriented singular Reidemeister moves

The purpose of this section is to give a generating set of oriented singular Reidemeister moves. We used this generating to write the axiom of our singquandle structure in section 3.

Minimal generating sets for oriented Reidemeister moves have been studied previously by Polyak in [15], where he proved that 4 moves are sufficient to generate the set for classical knots. However, analogous work for singular knot theory seems to be missing from the literature. In this section we will give a generating set of Reidemeister Moves for oriented singular links.

It was proven in [15] that the moves in Figure 5 constitute a minimal generating set of Reidemeister moves on oriented classical knots. For convenience we will use the same Ω notation used by Polyak [15] in our paper.



FIGURE 8. A generating set of Reidemeister moves for oriented knot.

To obtain a generating set of oriented singular Reidemeister moves we enumerate all possible such moves we then we show that they all can be obtained by a finite sequence of the moves given in Figure 2. Recall first that the nonoriented case of singular knots, we have the four moves given in Figure 5 singular Reidemeister moves.

If we consider the oriented case, we can easily compute the maximum number of Reidemeister moves to consider. For $\Omega 4$ moves we have 8 independent orientations, and for $\Omega 5$ we will have 6.



FIGURE 9. All four possible moves on nonoriented singular knots that involve singular crossings.

While a combinatorically driven approach yields a handful more, it is easily seen that some moves are simply the rotation of other moves, thus they are not considered. The 14 for Ω4 and Ω5 moves are tabulated in Figure 5 for reference.



FIGURE 10. All 14 oriented moves involving singular crossings.

Theorem 5.1. Only 3 oriented singular Reidemeister moves are required to generate the entire set of moves. These four moves are $\Omega 4a$, $\Omega 4e$, and $\Omega 5a$.

To show each moves dependence on this generating set, we will need to invoke Reidemeister moves of type $\Omega 1$ and $\Omega 2$ while performing transformations. The specific moves are given in [15]. To prove this theorem, we will first show that the moves of type $\Omega 4$ are generated by two unique moves. We will formulate this as a separate theorem:

Theorem 5.2. Only 2 oriented singular Reidemeister moves of type $\Omega 4$ are required to generate all type $\Omega 4$ moves. These moves are $\Omega 4a$ and $\Omega 4e$.

Lemma 5.3. The move $\Omega 4c$ is equivalent to $\Omega 2a \bigcup \Omega 4a \bigcup \Omega 2d$.

Proof.



□

Lemma 5.4. the move $\Omega 4d$ is equivalent to $\Omega 2c \bigcup \Omega 4a \bigcup \Omega 2d$.

Proof.



□

Lemma 5.5. the move $\Omega 4g$ is equivalent to $\Omega 2c \bigcup \Omega 4e \bigcup \Omega 2d$.

Proof.



□

Lemma 5.6. the move $\Omega 4h$ is equivalent to $\Omega 2c \bigcup \Omega 4e \bigcup \Omega 2d$.

Proof.



□

Lemma 5.7. the move $\Omega 4b$ is equivalent to $\Omega 2a \bigcup \Omega 2c \bigcup \Omega 4a \bigcup \Omega 2d \bigcup \Omega 2b$.

Proof.



□

By applying Lemma 5.4 we see that $\Omega 4d$ reduces further, and the lemma follows.

Lemma 5.8. the move $\Omega 4f$ is equivalent to $\Omega 2b \bigcup \Omega 2c \bigcup \Omega 4e \bigcup \Omega 2d \bigcup \Omega 2a$.

Proof.



□

By applying Lemma 5.6 we see that $\Omega 4h$ reduces further, and the lemma follows.

From here it remains to show that all moves of type $\Omega 5$ can be generated using only the $\Omega 5d$ move. We formulate this as a theorem:

Theorem 5.9. Only 1 oriented singular Reidemeister move of type $\Omega 5$ is required to generate all type $\Omega 5$ moves.

The proof of Theorem 5.9 is a consequence of the following Lemmas.

Lemma 5.10. The move $\Omega 5b$ is equivalent to $\Omega 1a \bigcup \Omega 4a \bigcup \Omega 5d \bigcup \Omega 4e \bigcup \Omega 1a$.

Proof.



□

Lemma 5.11. The move $\Omega 5c$ is equivalent to $\Omega 1b \bigcup \Omega 4a \bigcup \Omega 5d \bigcup \Omega 4e \bigcup \Omega 1b$.

Proof.



□

Lemma 5.12. The move $\Omega 5e$ is equivalent to $\Omega 1c \bigcup \Omega 4e \bigcup \Omega 5a \bigcup \Omega 4a \bigcup \Omega 1c$.

Proof.



□

Lemma 5.13. The move $\Omega 5f$ is equivalent to $\Omega 1d \bigcup \Omega 4e \bigcup \Omega 5a \bigcup \Omega 4a \bigcup \Omega 1d$.

Proof.



□

At this point, all $\Omega 5$ moves have been shown to depend on only $\Omega 4a$, $\Omega 4e$, $\Omega 5a$, and $\Omega 5d$. The last step remaining is to eliminate $\Omega 5d$.

Lemma 5.14. The move $\Omega 5d$ can be realized by a combination of $\Omega 2$ moves, and one $\Omega 5a$ move.

Proof.



□

6. Open questions

The following are some open questions for future research:

- Find other generating sets of oriented singular Reidemeister moves and prove their minimality.
- Define a notion of *extensions* of oriented singquandles as in [5].
- Define a cohomology theory of oriented singquandles and use low dimensional cocycles to construct invariants that generalize the number of colorings of singular knots by oriented singquandles.

References

[1] John C. Baez, *Link invariants of finite type and perturbation theory*, Lett. Math. Phys. **26** (1992), no. 1, 43–51, doi: 10.1007/BF00420517. MR1193625 (93k:57006)

[2] Khaled Bataineh, Mohamed Elhamdadi, and Mustafa Hajij, *The colored Jones polynomial of singular knots*, New York J. Math **22** (2016), 1439–1456.

[3] Joan S. Birman, *New points of view in knot theory*, Bull. Amer. Math. Soc. (N.S.) **28** (1993), no. 2, 253–287. MR1191478 (94b:57007)

[4] Joan S. Birman and Xiao-Song Lin, *Knot polynomials and Vassiliev's invariants*, Invent. Math. **111** (1993), no. 2, 225–270, doi: 10.1007/BF01231287. MR1198809

[5] J. Scott Carter, Mohamed Elhamdadi, Marina Appiou Nikiforou, and Masahico Saito, *Extensions of quandles and cocycle knot invariants*, J. Knot Theory Ramifications **12** (2003), no. 6, 725–738, doi: 10.1142/S0218216503002718. MR2008876

[6] Indu R. U. Churchill, Mohamed Elhamdadi, Mustafa Hajij, and Sam Nelson, *Singular Knots and Involutive Quandles*, arXiv:1608.08163, 2016.

[7] Mohamed Elhamdadi and Sam Nelson, *Quandles—an introduction to the algebra of knots*, Student Mathematical Library, vol. 74, American Mathematical Society, Providence, RI, 2015. MR3379534

[8] Mohamed Elhamdadi, Jennifer Macquarrie, and Ricardo Restrepo, *Automorphism groups of quandles*, J. Algebra Appl. **11** (2012), no. 1, 1250008, 9, doi: 10.1142/S0219498812500089. MR2900878

[9] Thomas Fiedler, *The Jones and Alexander polynomials for singular links*, J. Knot Theory Ramifications **19** (2010), no. 7, 859–866. MR2673687 (2012b:57024)

[10] V. F. R. Jones, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) **126** (1987), no. 2, 335–388, doi: 10.2307/1971403. MR908150

[11] David Joyce, *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Algebra **23** (1982), no. 1, 37–65, doi: 10.1016/0022-4049(82)90077-9. MR638121

[12] Jesús Juyumaya and Sofia Lambropoulou, *An invariant for singular knots*, Journal of Knot Theory and Its Ramifications **18** (2009), no. 06, 825–840.

[13] S. V. Matveev, *Distributive groupoids in knot theory*, Mat. Sb. (N.S.) **119(161)** (1982), no. 1, 78–88, 160 (Russian). MR672410

[14] Luis Paris, *The proof of Birman's conjecture on singular braid monoids*, Geom. Topol. **8** (2004), 1281–1300 (electronic). MR2087084

[15] Michael Polyak, *Minimal generating sets of Reidemeister moves*, Quantum Topol. **1** (2010), no. 4, 399–411, doi: 10.4171/QT/10. MR2733246

[16] V. A. Vassiliev, *Cohomology of knot spaces*, Theory of singularities and its applications, Adv. Soviet Math., vol. 1, Amer. Math. Soc., Providence, RI, 1990, pp. 23–69. MR1089670

Jordan University of Science and Technology, Irbid, Jordan
E-mail address: khaledb@just.edu.jo

University of South Florida, Tampa, USA
E-mail address: emohamed@mail.usf.edu

University of South Florida, Tampa, USA
E-mail address: mhajij@usf.edu

University of South Florida, Tampa, USA
E-mail address: wyoumans@mail.usf.edu